

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Characterization of Dermatoglyphics in PHOX2B-Confirmed Congenital Central Hypoventilation Syndrome

Emily S. Todd, Nicole M. Scott, Debra E. Weese-Mayer, Seth M. Weinberg,
Elizabeth M. Berry-Kravis, Jean M. Silvestri, Anna S. Kenny, Susan A. Hauptman,
Lili Zhou and Mary L. Marazita

Pediatrics 2006;118;408-414

DOI: 10.1542/peds.2005-3134

This information is current as of August 1, 2006

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://www.pediatrics.org/cgi/content/full/118/2/e408>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2006 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Characterization of Dermatoglyphics in *PHOX2B*-Confirmed Congenital Central Hypoventilation Syndrome

Emily S. Todd, MSPT^a, Nicole M. Scott, MPH^{b,c}, Debra E. Weese-Mayer, MD^a, Seth M. Weinberg, MA^b, Elizabeth M. Berry-Kravis, MD, PhD^{a,d}, Jean M. Silvestri, MD^a, Anna S. Kenny, CRT^a, Susan A. Hauptman, RRT^a, Lili Zhou, MD^a, Mary L. Marazita, PhD^{b,e,f}

Departments of ^aPediatrics and ^dNeurology and Biochemistry, Rush Children's Hospital at Rush University Medical Center, Chicago, Illinois; ^bCenter for Craniofacial and Dental Genetics, Division of Oral Biology and ^fDepartment of Oral and Maxillofacial Surgery, School of Dental Medicine and ^eDepartment of Human Genetics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania; ^cDepartment of Human Genetics, University of Michigan, Ann Arbor, Michigan

The authors have indicated they have no financial relationships relevant to this article to disclose.

ABSTRACT

OBJECTIVE. Individuals with congenital central hypoventilation syndrome have characteristic variants in the *PHOX2B* gene (primarily polyalanine expansion mutations). The *PHOX2B* gene acts as a transcriptional activator in the promotion of pan-neuronal differentiation in the autonomic nervous system during early embryologic development, with a primary role in the sympathetic noradrenergic phenotype in vertebrates. Because sympathetic innervation has been hypothesized to affect the development of dermatoglyphic pattern types, we hypothesized that individuals with *PHOX2B*-confirmed congenital central hypoventilation syndrome would have characteristic dermatoglyphic patterning and that the dermatoglyphic phenotype would be related to the disease-defining *PHOX2B* genotype.

METHODS. Dermatoglyphic pattern type frequency, left/right symmetry, and genotype/phenotype correlation were assessed for 33 individuals with *PHOX2B*-confirmed congenital central hypoventilation syndrome and compared with published control data.

RESULTS. Dermatoglyphic pattern type frequencies were altered in congenital central hypoventilation syndrome cases versus controls. In particular, there was an increase of arches in females and ulnar loops in males, with the largest differences for the left hand and for individuals with both congenital central hypoventilation syndrome and Hirschsprung disease. Dissimilarity scores between the congenital central hypoventilation syndrome and congenital central hypoventilation syndrome + Hirschsprung disease cases were not significantly different, nor were dissimilarity scores between all of the female and all of the male cases. No significant association was found between the number of polyalanine repeats in the *PHOX2B* genotypic category and dermatoglyphic pattern frequencies in the congenital central hypoventilation syndrome study groups.

www.pediatrics.org/cgi/doi/10.1542/peds.2005-3134

doi:10.1542/peds.2005-3134

Key Words

dermatoglyphics, *PHOX2B* gene, Hirschsprung disease

Abbreviations

CCHS—congenital central hypoventilation syndrome

ANS—autonomic nervous system

HD—Hirschsprung disease

SIDS—sudden infant death syndrome

Accepted for publication Mar 31, 2006

Address correspondence to Debra E. Weese-Mayer, MD, Rush University, Pediatric Respiratory Medicine at Rush Children's Hospital, 1653 West Congress Parkway, Chicago, IL 60612. E-mail: Debra_E_Weese-Mayer@rush.net

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275). Copyright © 2006 by the American Academy of Pediatrics

CONCLUSIONS. These results represent the first report describing specific dermatoglyphic patterning in congenital central hypoventilation syndrome and suggest a relationship between *PHOX2B* and the expression of dermatoglyphic pattern types. An expanded congenital central hypoventilation syndrome data set to include the full spectrum of *PHOX2B* mutations is necessary to further delineate the role of *PHOX2B* in dermatoglyphic patterning.

CHILDREN WITH CONGENITAL central hypoventilation syndrome (CCHS) characteristically present in the neonatal period with alveolar hypoventilation “in the absence of primary neuromuscular, lung, or cardiac disease, or an identifiable brainstem lesion.”¹ They demonstrate attenuated/absent responses to hypoxemia and hypercarbia, and related evidence of autonomic nervous system (ANS) dysfunction/dysregulation that may include Hirschsprung disease (HD) and/or tumors of neural crest origin.¹ Children with CCHS are heterozygous for a polyalanine expansion mutation in the paired-like homeobox (*PHOX2B*) gene,²⁻⁷ although a subset of patients has unique mutations in *PHOX2B*. Recently, a characteristic facial phenotype in CCHS has been described,⁸ with a pattern compatible with the developmental expression of *PHOX2B*. Long thought to be a rare disorder, CCHS is being diagnosed with increased frequency, since genetic testing for the *PHOX2B* mutation became clinically available in 2003.² As a result, general pediatricians may be caring for affected children in their practice.

The *PHOX2B* gene encodes a highly conserved homeobox domain transcription factor³ with an early embryologic action as a transcriptional activator in the promotion of pan-neuronal differentiation in the ANS and a separate role wherein it represses expression of inhibitors of neurogenesis.⁹⁻¹³ *PHOX2B* is considered to regulate the sympathetic noradrenergic phenotype in vertebrates.^{10,14} Sympathetic innervation has been hypothesized to affect the development of dermatoglyphic pattern types in primates, including humans,^{15,16} with dermatoglyphics defined as the study of the morphology and distribution of ridged skin patterns on the palmar surface of the hands and feet. We sought to more fully characterize the phenotype of CCHS by evaluating dermatoglyphics, an accessible phenotypic measure of embryologic development. We hypothesized that individuals with *PHOX2B*-confirmed CCHS would have characteristic dermatoglyphic patterning and that these pattern differences would vary with the clinical severity of CCHS (characterized by the presence or absence of HD) and the disease-defining *PHOX2B* genotype.

METHODS

Study Subjects

Diagnosis of CCHS was based on the American Thoracic Society Statement on CCHS¹ and confirmed with *PHOX2B* genotyping² (patent pending). Dermatoglyphic prints from 33 white individuals with *PHOX2B*-confirmed CCHS (all heterozygous for a polyalanine expansion mutation in *PHOX2B*) were analyzed, including 28 cases of isolated CCHS (14 females) and 5 cases with both CCHS and HD (2 females). CCHS subject age range was 3.0 to 20.75 years (mean: 13.6 ± 4.9 years). *PHOX2B* expansion mutations ranged from 25–33 repeats (mean: 27 ± 2 repeats). Control data were obtained from a published data set¹⁷ including 720 white children aged 7 years (360 females). This control data set was designed with the intent to provide a comparison sample for investigators studying dermatoglyphics in individuals with disease or congenital anomalies. Because fingerprints do not change during postnatal life, ages did not need to be matched. The study was approved by Rush University Medical Center and University of Pittsburgh institutional review boards. Informed consent and Health Insurance Portability and Accountability Act authorization were obtained from subjects and/or their parents or legal guardians.

Printing Technique and Dermatoglyphic Protocol

Fingerprints were obtained using a standard inkless technique¹⁸ using INFA-PRINT Inkless Imprint Wipes (Kansas City, MO). Three trained raters (E. S. T., N. M. S., D. E. W.-M.) independently categorized the dermatoglyphic prints as arch, ulnar loop, radial loop, or whorl.¹⁹ Figure 1 provides examples of each of these print types. Data sets were then merged, and any disagreements were reevaluated. Determination was made of each pattern frequency by specific study group (expressed as percentage of total digits evaluated in the group), by identification of subjects who are arch positive (≥ 1 arch on either hand), and by identification of subjects with ≥ 4 arches. These 3 determinations were conducted to allow for comparison to the control group data, as well as to the literature.

Statistical Analysis

χ^2 analysis was used to compare pattern frequencies for cases versus published control data,¹⁷ grouped by CCHS



FIGURE 1

This figure provides representative examples of dermatoglyphic patterns. These are from a right hand with the radial side of each print being to the left and the ulnar side to the right. A, Arch; B, Ulnar loop; C, Radial loop; D, Whorl.

status (all CCHS, isolated CCHS, or CCHS + HD), then stratified for gender and hand (left or right). Asymmetry was measured by dissimilarity scores, which were calculated by comparing prints on the paired homologous fingers of the left and right hands (eg, left index finger compared with right index finger). Paired fingers with dissimilar dermatoglyphic pattern types were given a score of 1, whereas paired fingers with the same pattern types were given a score of 0. Scores were then summed across all 5 finger pairs, thus, dissimilarity scores may range from 0 (having no dissimilarity) to 5 (having all 5 pairs of fingers dissimilar). The results were compared using the Wilcoxon-Mann-Whitney test. In addition, *PHOX2B* repeat number was compared with frequency of pattern type using linear regression with the continuous repeat number and logistic regression with repeat number divided into 2 categories: <29 repeats and ≥ 30 repeats. Dissimilarity scores were also compared between the 2 categories of *PHOX2B* repeat number using a Wilcoxon-Mann-Whitney test.²⁰ Results were considered significant if $P < .05$.

RESULTS

Dermatoglyphic Results

Pattern frequencies in CCHS cases and controls are provided in Table 1. Pattern frequencies in the text and table are shown as percentages of the total number of digits evaluated in each group. Gender differences in pattern frequency for cases, as well as arch positivity results, are provided below. Control values for arch positivity/multiple arches are unavailable, thus, no comparisons to controls could be made.

All CCHS Cases

Female cases had a higher arch frequency versus male cases (10.8 vs 3.6; $P = .01$) and a higher left-hand arch frequency versus male cases (14.1 vs 3.6; $P = .02$). Arch positivity (the presence of ≥ 1 digital arch) was seen in 30% of cases, with 60% of the arch-positive cases being female. Multiple arches (≥ 4 , no case had 3 arches) were identified in 12.1% of cases, with 75% of these being female.

Isolated CCHS

Arch positivity was seen in 25% of cases, with 57.1% of the arch-positive cases being female. Multiple arches were identified in 7.1% of cases, with 50% of these being female.

CCHS+HD

Female cases had a higher arch frequency versus male cases (both hands: 40.0 vs 0, $P = .002$; left hand: 40.0 vs 0, $P = .025$; right hand: 40.0 vs 0, $P = .025$). Arch positivity was seen in 60% of cases, with 66.7% of the arch-positive cases being female. Multiple arches were

identified in 40% of cases, with 100% of these being female.

Dissimilarity Scores

Dissimilarity scores between the CCHS and CCHS + HD cases were not different (1.11 and 0.75, respectively), nor were dissimilarity scores between all of the female cases and all of the male cases (mean: 1.0 and 1.13, respectively). Control values for dissimilarity scores are unavailable, thus, no comparisons to controls could be made.

Genotype/Phenotype Correlation

No significant association was found by linear regression or by logistic regression between the *PHOX2B* genotypic category (<29 polyalanine repeats versus ≥ 30 polyalanine repeats on the affected allele) and dermatoglyphic pattern frequencies in the CCHS study groups (all CCHS cases, isolated CCHS, and CCHS + HD).

DISCUSSION

These results represent the first report describing dermatoglyphics in individuals with *PHOX2B*-confirmed CCHS. Pattern type frequencies were altered in CCHS cases versus controls. In particular, there was an increase in arch and ulnar loop frequencies (typically seen in female and male cases, respectively), with the largest differences seen in the left hand and in the individuals with CCHS + HD. Dissimilarity scores as a proxy for asymmetry did not differ among the CCHS study groups or by *PHOX2B* genotypic category (<29 polyalanine repeats or ≥ 30 polyalanine repeats). Likewise, no significant association was found between the *PHOX2B* genotypic category and dermatoglyphic pattern frequencies.

Dermatoglyphics have been studied in children with isolated constipation and isolated HD,²¹⁻²⁵ both symptoms compatible with ANS dysregulation/dysfunction and both often reported among children with CCHS.²⁶ Gottlieb and Schuster²¹ reported arch positivity in 53% of patients and ≥ 4 arches in 27% of patients with early onset constipation (<10 years of age) (vs 13% and 2% of controls, respectively). Staiano et al²² demonstrated arch positivity in 38% of patients with constipation (vs 11% of controls) but no relationship between HD and digital arches. Drongowski and Coran²³ identified arch positivity in 33% of cases with functional chronic constipation, 32% of cases with anatomic chronic constipation (including HD), and 30% of control cases with inguinal hernia. Jackson et al²⁴ demonstrated arch positivity in 13% of constipated cases (vs 7% of controls). Goshima et al²⁵ demonstrated arch positivity in 28% of constipated patients (vs 24% of controls). The published results describing arch positivity in constipated patients are in keeping with our current results in children with CCHS. The published results describing arch positivity in patients with HD, however, are distinctly different from

TABLE 1 Digital Pattern Types and Laterality With CCHS Phenotype

Case	Pattern Types (%)				
	Hand	Arch	Ulnar Loop	Radial Loop	Whorl
All CCHS Cases					
Female	Left	14.1 ^c	55.1	3.8	26.9
	Right	7.6	64.6	0	27.8
	Both	10.8 ^a	59.9	1.9	27.4
Male	Left	3.6	69.9 ^c	3.6	22.9
	Right	3.6	63.9	2.4	30.1
	Both	3.6	66.9	3.0	26.5
Total	Left	8.7 ^a	62.7	3.7	24.8
	Right	5.6	64.2	1.2	29.0
	Both	7.1	63.5	2.5	26.9
Isolated CCHS					
Female	Left	10.3	54.4	4.4	30.9
	Right	3.0	65.2	0	31.9
	Both	6.6	59.9	2.2	31.4
Male	Left	4.1	69.9 ^b	1.4	24.7
	Right	4.1	61.6	1.4	32.9
	Both	4.1	65.8	1.4	28.8
Total	Left	7.1	62.4	2.8	27.7
	Right	3.5	63.4	0.7	32.4
	Both	5.3	62.9	1.8	30.0
CCHS + HD					
Female	Left	40.0 ^c	60.0	0	0
	Right	40.0 ^c	60.0	0	0
	Both	40.0 ^c	60.0 ^a	0	0
Male	Left	0	70.0	20.0	10.0
	Right	0	80.0	10.0	10.0
	Both	0	75.0 ^b	15.0 ^a	10.0
Total	Left	20.0 ^c	65.0	10.0	15.0
	Right	20.0 ^c	70.0	5.0	5.0
	Both	20.0 ^c	67.5	7.5	5.0 ^b
Controls					
Female	Left	4.6	64.9	3.4	27.2
	Right	6.2	64.2	4.0	25.5
	Both	5.4	64.5	3.7	26.4
Male	Left	4.2	64.7	5.4	25.7
	Right	3.8	59.1	5.0	32.0
	Both	4.0	62.0	5.2	28.8
Total	Left	4.4	64.8	4.4	26.5
	Right	5.0	61.7	4.5	28.8
	Both	4.7	63.3	4.5	27.6

P values are based on a Bonferroni correction and are a case/control comparison of frequencies. Total indicates female and male cases combined; Both, left and right hand combined.

^a *P* ≤ .01.

^b *P* < .001.

^c *P* < .0001.

our findings in children with CCHS + HD. These differences may be because of an intrinsic difference in the etiology and genetic basis of idiopathic HD versus syndromic HD.²⁷ Taken together, these findings extend the understanding of a relationship between dermatoglyphic patterning and functional/anatomic bowel disease, conditions that fall within the rubric of ANS dysregulation and will be encountered by the general care pediatrician.

The embryologic timing of the ridge pattern formation, between weeks 10 and 24 postfertilization, is associated with the type of dermatoglyphic pattern development. Early ridge formation is associated with whorls, late formation with arches, and intermediate formation

with loops. Clinical syndromes that have arrested embryologic development and decreased developmental maturation tend to have more arches and fewer whorls.²⁸ For example, an increase in arch and ulnar loop frequency and a decrease in whorl frequency have been reported in nonsyndromic female Filipinos with cleft lip and/or cleft palate.²⁹ Likewise, an increased incidence of arches^{30,31} and cleft lip and/or cleft palate³² has been reported in fetal hydantoin syndrome. Interestingly, female white sudden infant death syndrome (SIDS) victims have an increased arch frequency compared with the same control population we used.³³ Furthermore, male SIDS infants with dysmorphic features

or anatomic anomalies presented with more ulnar loops and fewer whorls, not dissimilar to the most affected cases with CCHS + HD in our study. Recognizing that children with CCHS have ANS dysregulation and that infants who have succumbed to SIDS also have ANS dysregulation,³⁴⁻⁴² the finding of altered dermatoglyphic patterning in CCHS and SIDS supports a relationship between the developing systems that regulate dermatoglyphics and altered embryologic development of the ANS. These interrelationships indicate that the study of dermatoglyphics provides an accessible means by which the pediatrician can probe the origin of complex diseases.

Although dermatoglyphic formation has been associated with many factors, including skeletal growth,²⁷ and volar pad topography,^{43,44} most interesting for the current study is that dermatoglyphic fingerprint patterns have been associated with the sympathetic innervation of the hand.^{15,16} We anticipate that the *PHOX2B* gene, as a regulator of the sympathetic noradrenergic phenotype in vertebrates,^{10,14} has a role in the determination of dermatoglyphic patterning, just as it has a role in the embryologic development of the enteric nervous system.^{10,45,46} This possibility is supported by our previous observation that 43% of children with CCHS have severe constipation, 21% of children with CCHS have HD,²⁶ and our current finding of an increased arch and ulnar loop frequency that is highest among children with *PHOX2B*-confirmed CCHS + HD. Our results here provide tentative support for the hypothesis that polyalanine expansion mutations in *PHOX2B* impact dermatoglyphic pattern formation, in addition to gut motility.

Also of interest is the noted gender difference in the pattern frequency analysis. These differences are notable in light of the well-recognized male predominance of isolated HD.⁴⁷ Recently, a common noncoding *RET* variant mutation within a conserved enhancer-like sequence has been identified as a contributing factor to the differences in susceptibility to HD by gender.⁴⁸ Most likely, the gender differences in frequency distribution of dermatoglyphics in CCHS are a result of the multifactorial inheritance of dermatoglyphic patterns, similar to HD. This is a more likely explanation for gender variation than a direct influence of the *PHOX2B* mutation, because we are unaware of other identified gender differences in the CCHS phenotype.

We have identified 2 potential limitations of this study. First, we relied on the published control data set¹⁷ including 720 white children aged 7 years (360 females) rather than recruit individual age-matched controls. This technique was used because it represents a better approximation of the population norms than a smaller set of matched controls. Furthermore, dermatoglyphic patterns do not change during an individual's postnatal life, unlike other phenotypic features, such as facial charac-

teristics, which have the potential to change with increasing age. Because of the data set used, we were not able to compare ≥ 1 or ≥ 3 to 4 arches with controls, which would have been useful in comparing our results with existing literature. Likewise, dermatoglyphic raters were not blinded to case versus control group, although it should be noted that the raters were blinded within CCHS clinical grouping. The second limitation is the small sample size of CCHS subjects. This may have reduced the ability to generalize our findings, but also may have contributed to the lack of significance when evaluated by *PHOX2B* genotype. In particular, the limited number of cases with higher numbers of polyalanine repeats (only 5 cases with 30 to 33 repeats) may have precluded detection of a significant correlation between the number of polyalanine repeats and frequencies of dermatoglyphic patterns.

CONCLUSIONS

Dermatoglyphic pattern frequency is clearly altered in CCHS cases versus controls, but the role of *PHOX2B* and the implication of the frequency differences are still to be determined. Based on these results, an increased sample size, including children with *PHOX2B*-confirmed CCHS unique mutations and a broad array of the common polyalanine expansion mutations, should be studied to fully understand the genotype/phenotype correlation in CCHS. Children with other disorders of neural crest origin, such as HD and neuroblastoma, should be studied to elucidate the relationship between these other disorders and an accessible phenotypic measure of the embryological neural crest. It would also be worthwhile to evaluate dermatoglyphics of parents and siblings of individuals with CCHS to look for an absence of familial tendencies, thus supporting a causative effect of the *PHOX2B* mutation on the dermatoglyphic pattern frequencies. In so doing, the dermatoglyphic phenotype of CCHS will be further characterized, and the contribution of *PHOX2B* to both dermatoglyphic patterning and CCHS will be better understood.

ACKNOWLEDGMENTS

This work was supported by Rush University Dean's Office Summer Research Fellowship and Sigma Xi Outstanding Rush Medical College Student Research Award (E. S. Todd), Spastic Paralysis and Allied Diseases of the Central Nervous System Research Foundation of the Illinois-Eastern Iowa District Kiwanis International (E. M. Berry-Kravis), and National Institutes of Health/National Institute of Dental and Craniofacial Research grants R01-DE016148 and P50-DE016215 (M. L. Marazita).

We thank Kelly Dearing Roollen and Indentiseal/Infra-Print for their gracious donation of the inkless fingerprinting materials to make this study possible. We also thank Foster Lewin for the preparation of the DNA,

Chinton Desai and Ronak Lahkia who assisted with PCR assays, and the children with CCHS who graciously shared their DNA and their fingerprints.

REFERENCES

1. Weese-Mayer DE, Shannon DC, Keens TG, Silvestri JM. American Thoracic Society statement on the diagnosis and management of idiopathic congenital central hypoventilation syndrome. *Am J Respir Crit Care Med.* 1999;160:368–373
2. Weese-Mayer DE, Berry-Kravis EM, Zhou L, et al. Idiopathic congenital central hypoventilation syndrome: Analysis of genes pertinent to early autonomic nervous system embryologic development and identification of mutations in *PHOX2B*. *Am J Med Gen.* 2003;123A:267–278
3. Amiel J, Laudier B, Attie-Bitach T, et al. Polyalanine expansion and frameshift mutations of the paired-like homeobox gene *PHOX2B* in congenital central hypoventilation syndrome. *Nat Genet.* 2003;33:459–461
4. Matera I, Bachetti T, Puppo F, et al. *PHOX2B* mutations and polyalanine expansions correlate with the severity of the respiratory phenotype and associated symptoms in both congenital and late onset central hypoventilation syndrome. *J Med Gen.* 2004;41:373–380
5. Sasaki A, Kanai M, Kijima K, et al. Molecular analysis of congenital central hypoventilation syndrome. *Hum Genet.* 2003;114:22–26
6. Trang H, Dehan M, Beaufile F, Zaccaria I, Amiel J, Gaultier C. The French Congenital Central Hypoventilation Syndrome registry: general data, phenotype, and genotype. *Chest.* 2005;127:72–79
7. Trochet D, O'Brien LM, Gozal D, et al. *PHOX2B* genotype allows for prediction of tumor risk in congenital central hypoventilation syndrome. *Am J Hum Genet.* 2005;76:421–426
8. Todd ES, Weinberg SM, Berry-Kravis EM, et al. Facial phenotype in children and young adults with *PHOX2B*-determined congenital central hypoventilation syndrome: Quantitative pattern of dysmorphology. *Pediatr Res.* 2006;59:39–45
9. Dubreuil V, Hirsch M, Jouve C, Brunet J, Goridis C. The role of *PHOX2b* in synchronizing pan-neuronal and type-specific aspects of neurogenesis. *Development.* 2002;129:5241–5253
10. Pattyn A, Morin X, Cremer H, Goridis C, Brunet JF. The homeobox gene *Phox2b* is essential for the development of autonomic neural crest derivatives. *Nature.* 1999;399:366–370
11. Stanke M, Junghans D, Geissen M, Goridis C, Ernsberger U, Rohrer H. The Phox2 homeodomain proteins are sufficient to promote the development of sympathetic neurons. *Development.* 1999;126:4087–4094
12. Samad OA, Geisen MJ, Caronia G, et al. Integration of antero-posterior and dorsoventral regulation of *Phox2b* transcription in cranial motoneuron progenitors by homeodomain proteins. *Development.* 2004;131:4071–4083
13. Huber K, Karch N, Ernsberger U, Goridis C Unsicker K. The role of *Phox2B* in chromaffin cell development. *Dev Biol.* 2005;279:501–508
14. Lo L, Morin X, Brunet J, Anderson DJ. Specification of neurotransmitter identity by *Phox2* proteins in neural crest stem cells. *Neuron.* 1999;22:693–705
15. Dell DA, Munger BL. The early embryogenesis of papillary (sweat duct) ridges in primate glabrous skin: the dermatotopic map of cutaneous mechanoreceptors and dermatoglyphics. *J Comp Neurol.* 1986;244:511–532
16. Moore SJ, Munger BL. The early ontogeny of the afferent nerves and papillary ridges in human digital glabrous skin. *Brain Res Dev Brain Res.* 1989;48:119–141
17. Plato CC, Cereghino JJ, Steinberg SS. Dermatoglyphics of American Caucasians. *Am J Phys Anthropol.* 1975;42:195–210
18. Walker NF. Inkless methods of finger, palm, and sole printing. *J Pediatr.* 1957;50:27–29
19. Cummins H, Mildo WC. *Fingerprints, Palms and Soles: An Introduction to Dermatoglyphics*. New York, NY: Dover; 1961
20. SAS Institute Inc. *SAS Procedures Guide*. Version 8. Cary, NC: SAS Institute Inc; 1999
21. Gottlieb SH, Schuster MM. Dermatoglyphic (fingerprint) evidence for a congenital syndrome of early onset constipation and abdominal pain. *Gastroenterology.* 1986;91:428–432
22. Staiano A, Andreotti MR, Perrotta V, Strisciuglio P. Prevalence of digital arches in children with abdominal pain and constipation. *J Pediatr.* 1990;117:435–436
23. Drongowski RA, Coran AG. Dermatoglyphic patterns in children with chronic constipation. *Dig Dis Sci.* 1995;40:1420–1422
24. Jackson CR, Anderson B, Jaffray B. Childhood constipation is not associated with characteristic fingerprint patterns. *Arch Dis Child.* 2003;88:1076–1077
25. Goshima S, Fagundes-Neto U, Morais MB. Dermatoglyphics of children with chronic constipation. *Arq Gastroenterol.* 2004;41:37–41
26. Weese-Mayer DE, Silvestri JM, Huffman AD, et al. Case/control family study of ANS dysfunction in idiopathic congenital central hypoventilation syndrome. *Am J Med Genetics.* 2001;100:237–245
27. Brooks AS, Oostra BA, Hofstra RMW. Studying the genetics of Hirschsprung's disease: unraveling an oligogenic disorder. *Clin Genet.* 2004;67:6–14
28. Babler WJ. Embryologic development of epidermal ridges and their configurations. *Birth Defects Orig Artic Ser.* 1991;27:95–112
29. Scott NM, Weinberg SM, Neiswanger K, et al. Dermatoglyphic pattern types in subjects with nonsyndromic cleft lip with or without cleft palate (CL/P) and their unaffected relatives in the Philippines. *Cleft Palate Craniofac J.* 2005;42:362–366
30. Lu MC, Sammel MD, Cleveland RH, Ryan LM, Holmes LB. Digit effects produced by prenatal exposure to antiepileptic drugs. *Teratology.* 2000;61:277–283
31. Bokhari A, Coull BA, Holmes LB. Effect of prenatal exposure to anticonvulsant drugs on dermal ridge patterns of fingers. *Teratology.* 2002;66:19–23
32. Jones KL, ed. *Smith's Recognizable Patterns of Human Malformation*. 5th ed. Philadelphia, PA: W.B. Saunders Company; 1997: 559–563
33. Kozakewich H, Fox K, Plato CC, Cronk C, Mandell F, Vawter GF. Dermatoglyphics in sudden infant death syndrome. *Pediatr Pathol.* 1992;12:637–651
34. Franco P, Szliwowski H, Dramaix M, Kahn A. Decreased autonomic responses to obstructive sleep events in future victims of sudden infant death syndrome. *Pediatr Res.* 1999;46:33–39
35. Kelly DH, Golub H, Carley D, Shannon DC. Pneumograms in infants who subsequently died of sudden infant death syndrome. *J Pediatr.* 1986;109:249–254
36. Franco P, Szliwowski H, Dramaix M, Kahn A. Polysomnographic study of the autonomic nervous system in potential victims of sudden infant death syndrome. *Clin Auton Res.* 1998;8:243–249
37. Ledwidge M, Fox G, Matthews T. Neurocardiogenic syncope: a model for SIDS. *Arch Dis Child.* 1998;78:481–483
38. Schechtman VL, Harper RM, Kluge KA, Wilson AJ, Hoffman HJ, Southall DP. Cardiac and respiratory patterns in normal infants and victims of the sudden infant death syndrome. *Sleep.* 1988;11:413–424
39. Meny RG, Carroll JL, Carbone MT, Kelly DH. Cardiorespiratory recordings from infants dying suddenly and unexpectedly at home. *Pediatr.* 1994;93:44–49
40. Ponsonby AL, Dwyer T, Gibbons LE, Cochrane JA, Jones ME, McCall MJ. Thermal environment and sudden infant

- death syndrome: case-control study. *BMJ*. 1992;304:277-282
41. Kahn A, Groswasser J, Rebuffat E, et al. Sleep and cardiorespiratory characteristics of infant victims of sudden death: a prospective case-control study. *Sleep*. 1992;15:287-292
 42. Schwartz PJ, Stramba-Badiale M, Segantini A, et al. Prolongation of the QT interval and the sudden infant death syndrome. *N Engl J Med*. 1998;338:1709-1714
 43. Bonnevie K. Studies on papillary patterns of human fingers. *J Genet*. 1924;15:1-111
 44. Mulvihill JJ, Smith DW. The genesis of dermatoglyphics. *J Pediatr*. 1969;75:579-589
 45. Pattyn A, Morin X, Cremer H, Goridis C, Brunet J. Expression and interaction of the closely related homeobox genes *Phox2a* and *Phox2b* during neurogenesis. *Development*. 1997;124:4065-4075
 46. Young HM, Ciampoli D, Hsuan J, Carty AJ. Expression of Ret-p^{75NTR}, Phox2a-, Phox2b-, and tyrosine hydroxylase immunoreactivity by undifferentiated neural crest derived cells and different classes of enteric neurons in the embryonic mouse gut. *Dev Dyn*. 1999;216:137-152
 47. Badner JA, Sieber WK, Garver KL, Chakravarti A. A genetic study of Hirschsprung disease. *Am J Med Genet*. 1990;46:568-580
 48. Emison ES, McCallion AS, Kashuk, CS, et al. A common sex-dependent mutation in a *RET* enhancer underlies Hirschsprung disease risk. *Nature*. 2005;434:857-863

Characterization of Dermatoglyphics in PHOX2B-Confirmed Congenital Central Hypoventilation Syndrome

Emily S. Todd, Nicole M. Scott, Debra E. Weese-Mayer, Seth M. Weinberg,
Elizabeth M. Berry-Kravis, Jean M. Silvestri, Anna S. Kenny, Susan A. Hauptman,
Lili Zhou and Mary L. Marazita
Pediatrics 2006;118;408-414
DOI: 10.1542/peds.2005-3134

This information is current as of August 1, 2006

Updated Information & Services	including high-resolution figures, can be found at: http://www.pediatrics.org/cgi/content/full/118/2/e408
References	This article cites 45 articles, 11 of which you can access for free at: http://www.pediatrics.org/cgi/content/full/118/2/e408#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Genetics & Dysmorphology http://www.pediatrics.org/cgi/collection/genetics_and_dysmorphology
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.pediatrics.org/misc/Permissions.shtml
Reprints	Information about ordering reprints can be found online: http://www.pediatrics.org/misc/reprints.shtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

